Mon. Not. R. Astron. Soc. **000**, 1–5 (2010) Printed 20 September 2010 (MN LATEX style file v2.2)

# Boosting hierarchical structure formation with scalar-interacting dark matter

Wojciech A. Hellwing[1⋆], Steffen R. Knollmann[2] and Alexander Knebe[2]
[1]*Nicolaus Copernicus Astronomical Center, ul. Bartycka 18, Warsaw, 00-716, Poland*
[2]*Grupo de Astrofísica, Departamento de Física Teórica, Módulo C-15, Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Cantoblanco, Madrid, Spain*

arXiv:1004.2929v3 [astro-ph.CO] 17 Sep 2010

Accepted 2010 August 22 . Received 2010 August 11; in original form 2010 April 16

**ABSTRACT**
We investigate the effect of long-range scalar interactions in dark matter (DM) models of cosmic structure formation with a particular focus on the formation times of haloes. Utilising $N$-body simulations with $512^3$ DM particles we show that in our models dark matter haloes form substantially earlier: tracing objects up to redshift $z \sim 6$ we find that the formation time, as characterised by the redshift $z_{1/2}$ at which the halo has assembled half of its final mass, is gradually shifted from $z_{1/2} \approx 1.83$ in the fiducial $\Lambda$CDM model to $z_{1/2} \approx 2.54$ in the most extreme scalar-interaction model. This is accompanied by a shift of the redshift that marks the transition between merger and steady accretion epochs from $z_* \approx 4.32$ in the $\Lambda$CDM halos to $z_* \approx 6.39$ in our strongest interaction model. In other words, the scalar-interacting model employed in this work produces more structures at high redshifts, prolonging at the same time the steady accretion phases. These effects taken together can help the $\Lambda$CDM model to account for a high redshift reionisation as indicated by the WMAP data and can alleviate issues related to the survival of the thin-disk dominated galaxies at low redshifts.

**Key words:** methods: n-body simulations – galaxies: halos – galaxies: evolution – cosmology: theory – dark matter

## 1 INTRODUCTION

The $\Lambda$ cold dark matter ($\Lambda$CDM) model has proven to be capable of explaining a tremendous amount of observational data. As we entered the era of precision cosmology — both observationally and from a modellers perspective — we are left with a more and more detailed picture of what actually took place when the Universe cooled down and formed structures starting from $z \sim 1100$ to the present day. But there are still issues related to the structure formation on galactic scales to be resolved. For instance, observations of high redshift quasars (Willott et al. 2010) and galaxies (Ouchi et al. 2009) imply that the epoch of the reionisation of the Universe ended at $z \sim 6$. On the other hand observations of the polarisation of the cosmic microwave background radiation (Komatsu & et al. 2010; Benson et al. 2006; Kogut et al. 2003) suggest that the Universe was already reionised at $z \gtrsim 10 - 11$. However, galaxy formation within the $\Lambda$CDM model using the suggested moderate normalisation of the power spectrum of primordial density perturbations $\sigma_8 = 0.8$ (as favoured by recent WMAP observations; Komatsu & et al. 2010) may not happen early enough to account for such a high redshift of reionisation. Furthermore the astonishing array of recent observations of ultra-faint distant objects (Bouwens et al. 2010, 2009) as well as advancements in the understanding of the high-$z$ physics (González et al. 2010; Haiman 2004; Volonteri et al. 2003) have shown that the young Universe was a busy and already structure-rich place. In addition, there are issues related to the survival of thin-disk dominated galaxies. It was shown that in a pure CDM model of galaxy-halo formation, the majority of Milky-Way-sized haloes with $M_0 \sim 10^{12} h^{-1} M_\odot$ have accreted at least one object with greater total mass than the Milky Way disk ($m > 5 \times 10^{10} h^{-1} M_\odot$) over the redshifts $z \leqslant 3$ (Driver 2010; Stewart et al. 2008). It has been shown recently (Governato et al. 2009; Moster et al. 2010), that the disk may reform after mergers or even survive minor mergers, somewhat easing the problem of disk survival. Still, late-time bombardment of Milky-Way like haloes, predicted within the $\Lambda$CDM model, could still be considered as a concern about the survival of thin-disk-dominated galaxies and the existence of galaxies with unusually quiet merger history like our Galaxy. Moreover, recent observations of the distant Universe suggest that much of the stellar mass of bright galaxies was already present in place at $z > 1$, this presents the "downsizing" challenge for models of galaxy formation because massive haloes are assembled late in the hierarchical clustering process intrinsic to the CDM cosmology (Cowie et al. 2003; Fontanot et al. 2009; Bower et al. 2006). ~~Finally~~ In addition there were also reports on massive superclusters found at intermediate redshifts ($z \sim 1$), which seem to be very rare and extreme objects within standard $\Lambda$CDM model (Yamila Yaryura et al. 2010;

⋆ E-mail: pchela@camk.edu.pl

Colless & et al. 2003; Swinbank & et al. 2007). Finally there is a recent claim of possible observational evidence for a "fifth force" in DM sector (Lee 2010).

Even though we would not go as far as to call this a "galaxy formation crisis", there are noticeable tensions between what observations suggest and what (cosmological simulations of) the ΛCDM model seems to predict. The ReBEL (*daRk Breaking Equivalence principLe*) model discussed in recent years in the literature (Gubser & Peebles 2004b,a; Farrar & Peebles 2004; Farrar & Rosen 2007; Peebles & Nusser 2010) has the potential to alleviate these tensions and to help the theoretical ΛCDM model to better match the observations. The ReBEL model is a slight modification of the standard cold dark matter model. It introduces an additional long-range scalar mediated force only in the dark sector. This has been studied in numerical simulations (Nusser et al. 2005; Hellwing & Juszkiewicz 2009; Hellwing 2010; Keselman et al. 2010; Hellwing et al. 2010) and these studies have shown that the model can match observational properties of the large-scale structures (like the one-dimensional Ly-$\alpha$ power spectrum and the power spectrum of the SDSS[1] galaxies) as well as the standard ΛCDM does, providing at the same time interesting features with respects to the formation of galactic structures. We also acknowledge that recently other authors performed structure formation tests in similar scalar-interacting DM scenarios (Baldi et al. 2010; Baldi 2009; Li & Barrow 2010; Li & Zhao 2010), but they focused on models containing scalar-field-like Dark Energy interacting with DM. Elementary considerations related to the ReBEL model and the reionisation mechanism showed that scalar-interacting DM could accommodate early reionisation of the intergalactic medium (Cen 2006). The low-resolution simulations presented in the ReBEL literature have indicated that this model can foster a picture in which structure formation starts at higher redshifts. In this letter we will show that this is indeed the case. We will present, for the first time in the literature, results of analysis of the high-resolution N-body simulations of hierarchical DM halo formation in the ReBEL framework.

## 2 SELF-INTERACTING DM: THE REBEL MODEL

The theoretical ground of the ReBEL model was formed in two papers by Gubser & Peebles (2004b,a). We follow the phenomenological description of this scalar-interacting DM model presented in Nusser et al. (2005) and Hellwing & Juszkiewicz (2009). We consider a modified effective potential and force law between DM particles of the forms

$$\Phi(r) = -\frac{Gm}{r}\left(1+\beta e^{-r/r_s}\right), \quad (1)$$

$$F_{DM} = -\frac{Gm^2}{r^2}\left[1+\beta\left(1+\frac{r}{r_s}\right)e^{-r/r_s}\right], \quad (2)$$

where the two free parameters of the model are: $\beta$, the measure of the strength of the scalar force compared to usual Newtonian gravity, and $r_s$, the screening length of the additional force that is constant in a comoving frame. Previous considerations in the literature of the subject provide a crude estimate of the values of this free parameters to be of orders $\beta \sim 1$ and $r_s \sim 1h^{-1}\,\mathrm{Mpc}$. In this short report we explore two possible values of the $\beta = 0.5, 1$ and two values of the screening length parameter $r_s = 0.5, 1h^{-1}\,\mathrm{Mpc}$. We label the corresponding simulations runs as ΛCDM, b05rs05, b05rs1, b1rs1, where b stands for $\beta$ and rs marks the $r_s$ parameter. In the ΛCDM run we use $\beta = 0$ (i.e. no scalar forces). Equations (1,2) yield a simple phenomenological description of the ReBEL addition to the CDM model.

## 3 NUMERICAL MODELLING

To follow the formation of structures within the ReBEL framework we use an adapted version of the `GADGET2` code (Springel 2005). For the detailed descriptions of the modifications made to the code we refer the reader to our previous paper on this subject (Hellwing & Juszkiewicz 2009). We have conducted a series of high-resolutions DM only $N$-body simulations containing $512^3$ particles within periodic box of $32h^{-1}\,\mathrm{Mpc}$ comoving width. The cosmology used in the simulations was the canonical ΛCDM with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $\sigma_8 = 0.8$ and $h = 0.7$. Thus our mass resolution is $m_p \simeq 2.033\times10^7 h^{-1} M_\odot$. The force softening parameter was set to be $\varepsilon = 6h^{-1}\,\mathrm{kpc}$. All simulations were started with the same initial conditions at $z_{ini} = 50$, and varied only in the ReBEL parameters. For each run we saved 30 snapshot equally spaced in the logarithmic time scale starting from $z = 6.092$ to $z = 0$.

We used the MPI+OpenMP hybrid `AMIGA` halo finder (`AHF`) to identify haloes and subhaloes in our simulation[2]. `AHF` is the successor of the `MHF` halo finder by Gill et al. (2004), a detailed description of AHF is given in the code paper (Knollmann & Knebe 2009). We note that we have adjusted the code slightly to take the modified gravity of the ReBEL models into account. Starting with the halo catalogues at $z = 0$ we construct the merger trees for each simulation by cross-correlating the particles constituting the haloes in consecutive time steps. For each halo we record all progenitors and select the main progenitor as the halo that maximises the merit function $N_1N_2/N_s^2$, where $N_{1,2}$ and $N_s$ are the number of particles of the haloes and the number of shared particles, respectively.

## 4 THE RESULTS

In this *Letter* we focus primarily on the mass accretion histories of dark matter haloes and their masses, respectively. A more in-depth study of the effects of the ReBEL model on their internal properties will be presented in a companion paper (Hellwing, Knollman & Knebe, in preparation).

### 4.1 Mass accretion histories

Following the scheme presented in Wechsler et al. (2002) (hereafter W02), we fit each halo's mass accretion history (MAH) to the exponential law of the form

$$\tilde{M}(a) = \exp[\alpha(1-a^{-1})], \quad \tilde{M}(a) \equiv \frac{M(a)}{M_0}, \quad (3)$$

where $\tilde{M}$ is the halo mass for a given cosmic scale factor $a$ expressed in terms of its final mass $M_0$ at $z = 0$ and $\alpha$ is a free parameter to be determined via fitting.

The fit parameter can be used to express a characteristic redshift of formation:

$$z_* = \frac{S}{\alpha} - 1, \quad (4)$$

[1] Sloan Digital Sky Survey - `http://www.sdss.org/`

[2] AHF is freely available from `http://popia.ft.uam.es/AMIGA`




**Figure 1.** The probability distribution of the formation redshift $z_{1/2}$ for all haloes containing at least 500 particles ($M_0 > 10^{10}h^{-1}M_\odot$).

defined as the time when the logarithmic slope of the accretion rate $\mathrm{d}\log M/\mathrm{d}\log a$ falls below some specified value $S$. In our studies we adopt vale $S = 2$ advocated in W02. Therefore our $z_*$ is the redshift indicating the time when a dark matter halo has entered the steady accretion phase of its mass accumulation (i.e. the last major merger should have taken place at $z > z_*$). The formation redshift — usually defined as the time when an object constituted half of its present-day mass (e.g. Lacey & Cole 1993) — can now be expressed in terms of $z_*$

$$z_{1/2} = \frac{-\log 0.5}{S}(z_* + 1) = \frac{-\log 0.5}{\alpha}\,. \qquad (5)$$

We will use $z_*$ and $z_{1/2}$ as obtained by fitting our numerical MAHs to eqn. 3 as estimators of the steady accretion transitions and the formation times of our halos. We acknowledge that our derivation of both these redshifts is obviously not independent. To this extent, we also derived the formation redshift $z_{1/2}$ by solely using the numerical MAH of each halo finding the point where the mass drops below $M_0/2$; the cross-correlation between these two $z_{1/2}$ values clearly follows a straight 1:1 relation with a scatter of less than 5%. We therefore decided to stick to the value obtained by eqn. (5).

In general, eqn. (3) was shown to be a universal good fit to the halos MAH's (e.g. W02), however Tasitsiomi et al. (2004) pointed out that the simple exponential one parameter form of W02 is not a good fit to the MAH of cluster haloes as well as haloes that experienced recent major mergers. To not be biased by including such objects in our study we removed all haloes whose $\chi^2$-value deviated from the mean by more than $1\sigma$ and all haloes whose Most Massive Progenitor (MMP) was not present in at least 24 (out of 30) snapshots, thus eliminating of order $\sim 25$ per cent of all haloes in the original catalogues. We further limit the analysis to objects containing in excess of 500 particles (at redshift $z = 0$) corresponding to a lower mass cut of $\sim 10^{10}h^{-1}M_\odot$. This leave us with $\sim 7000$ objects in the ΛCDM sample, and $\sim 4700$ objects in the b1rs1 ReBEL run.

### 4.2 Formation and Characteristic Redshifts

When plotting the distribution function of formation times $z_{1/2}$ in Fig. 1 we observe a clear trend for the mean of $z_{1/2}$ to shift to higher redshifts in the ReBEL models. We like to note in passing that for the ΛCDM model our results agree with the ones presented

**Table 1.** The mean and the $1\sigma$ scatter of the $z_f$ and $z_*$ distributions.

| The model ($\beta/r_s$) | $\langle z_{1/2}\rangle$ | $\sigma_{z_{1/2}}$ | $\langle z_*\rangle$ | $\sigma_{z_*}$ |
|---|---|---|---|---|
| ΛCDM | 1.83 | 0.8 | 4.32 | 2.3 |
| $0.5/0.5h^{-1}$ Mpc | 2.12 | 0.9 | 5.16 | 2.51 |
| $0.5/1.0h^{-1}$ Mpc | 2.21 | 0.9 | 5.4 | 2.7 |
| $1.0/1.0h^{-1}$ Mpc | 2.54 | 1.0 | 6.39 | 3.0 |




**Figure 2.** Hierarchical structure formation for all models. The relation between the formation redshift ($z_{1/2}$) of objects and their masses $M$. The error-bars represents $1\sigma$ scatter within the binned distributions. The error-bars of the ReBEL model where slightly shifted to the right for clarity.

in Lin et al. (2003, their Fig. 2 and 3), when using the same mass bins as in said work.

We calculated the same kind of distribution for the characteristic redshift $z_*$ (though not explicitly shown here) and list for both distributions the respective means and variances in Tab. 1. We note that for the strongest ReBEL model considered in this *Letter* the mean value of $z_{1/2}$ is 39 per cent larger then the corresponding $\Lambda CDM$ value. This effect is even more pronounced for the mean values of the $z_*$ distributions, the corresponding shift is more than 48 per cent there. The corresponding $1\sigma$ scatters are also larger. This is not surprising as these two quantities are linked with linear relation of eqn. (5). We also note that the magnitude of changes of $\langle z_{1/2}\rangle$ and $\langle z_*\rangle$ in models *b05rs05* and *b05rs1* are of comparable order. The reason for this is the following: our sample is dominated by objects with masses $M_0 < 10^{12}h^{-1}M_\odot$ whose virial radii are $r_{vir} < 200h^{-1}$ kpc; therefore for majority of our objects the screening length of the ReBEL force is much greater than their radii, hence for this class of objects only the $\beta$ parameter measures overall enhancement of scalar-attraction. To conclude this paragraph we denote that for the parameter space probed by our simulations the $\beta$ parameter plays a more important role on the examined quantities than the screening length.

Viewing the shifts for formation and characteristic redshift in combination we determine that, on average, haloes in the ReBEL model form earlier, thus arriving at moderate and low redshifts with higher masses. Moreover haloes with exceptionally long quiet accretion epochs are much more abundant in the populations of our ReBEL models. We will return to the issue of larger masses below in Sec. 4.3 where we study the halo mass function at various redshifts.

To check whether the hierarchical structure formation scenario




**Figure 3.** The cumulative mass functions for all haloes containing 20 and more particles calculated at redshifts $z = 6, 3$ and 0. Mass functions for redshifts $z = 6$ ($z = 3$) were scaled down by $10^{-2}$ $(10^{-1})$ for clarity.

is preserved in the ReBEL models and how the effect of the earlier structure formation depends on the halo mass we show in Fig. 2 the average $z_{1/2}$ binned in halo mass. We see that the hierarchical character of the structure formation is preserved in all the simulations runs, i.e. small mass objects form first with higher mass objects forming later. In all the ReBEL runs the formation redshifts are higher for all considered masses, but the effect is strongest for less massive haloes. However, we also notice a clear dependence on the parameters of the ReBEL model. For small $\beta$ and small $r_s$ the effect of shifting structure formation to the higher redshifts is less pronounced. This is of course expected, since the screening length acts here as the effective radius of an enhanced accretion and the $\beta$ measure the strength of this enhancement. It is interesting to note, that for the *b05rs05, b05rs1* runs the aforementioned effect disappears at some fixed mass scale. In other words, we see that the averaged ReBEL $z_{1/2}$ values start to agree with the ΛCDM values starting from mass $M \sim 5 \times 10^{12} h^{-1} M_{\odot}$. It is only the *b1rs1* model that deviates throughout the whole mass range probed by our simulations. Nevertheless we observe that in the *b05rs05, b05rs1* simulations the haloes with masses $10^{11} h^{-1} M_{\odot} < M < 5 \times 10^{12} h^{-1} M_{\odot}$ form at significantly higher redshift compared to the standard CDM model. For the *b1rs1* case even haloes with masses comparable to a mass of a group of galaxies $10^{13} h^{-1} M_{\odot} < M < 10^{14} h^{-1} M_{\odot}$ acquire most of their mass at relatively earlier times, however we denote that at masses $M > 10^{13} h^{-1} M_{\odot}$ the signal is dominated by small-number effects, i.e. we only have of order $\sim 30$ objects in that mass range: to more thoroughly check the differences in ΛCDM and ReBEL models at this mass scale simulations with a larger box size are required.

### 4.3 The Evolution of the Mass Function

As speculated before, haloes in the ReBEL models not only form earlier but also carry a larger mass. To actually confirm and quantify this previously made claim (cf. Sec. 4.2) we plot in Fig. 3 the cumulative mass functions (CMFs) at the redshifts $z = 6, 3$, and 0.

We notice that for $z = 6$ and $z = 3$ all ReBEL's CMFs have higher amplitudes than the fiducial ΛCDM CMF in the whole probed mass range: At high redshifts the ReBEL model produces more non-linear structures compared to the standard CDM model. However structure formation in the ΛCDM paradigm seems to be 'catching' up with ReBEL at lower redshifts. This can again be seen in Fig. 3: the discrepancies of the amplitudes at redshift $z = 0$ are much suppressed in comparison to the high redshift CMF's. In addition, we note an interesting feature at the final redshift: while the CMFs have a higher amplitude in the ReBEL models for redshifts $z = 6$ and 3 across the whole mass range, at $z = 0$ only the high mass end of the ReBEL mass functions shows an excess relative to the fiducial ΛCDM model. The low mass end of the ReBEL CMFs at this redshift show deficiency of haloes compared to the LCDM. The latter effect indicates that in the ReBEL model some of the low mass haloes were already used in structure formation processes as bricks for more massive objects. All this taken together clearly points out that the process of rapid non-linear structure formation takes place at later times in the ΛCDM model compared to model with the ReBEL forces.

## 5 DISCUSSION

The main conclusion of this *Letter* is that a slight modification to the CDM model in the form of scalar interactions can accommodate more gravitationally bound structures at high redshifts (e.g. fig. 3) which in turn promotes an early reionisation of the Universe. This can be expressed by examining the $\eta$ factor that describes the efficiency of reionisation of the Inter Galactic Medium (IGM) (Cen 2006, 2003):

$$\eta \equiv \frac{c_{\star} f_{esc} (\mathrm{d}f_M/\mathrm{d}t) e_{UV}}{C(1+z)^3}\,, \tag{6}$$

where $c_{\star}$ is the star formation factor efficiency, $f_{esc}$ is the ionising photon escape fraction, $\mathrm{d}f_M/\mathrm{d}t$ is the halo formation rate, $e_{UV}$ is the ionising photon production efficiency and $C$ is the gas clumping factor. Here the numerator reflects the ionising photon production rate while the denominator is the ionising photon destruction rate. When $\eta$ exceeds a certain threshold, the IGM becomes ionised. As the ReBEL models exhibit a richer structure at $z = 6$, their formation rate must have been higher than the one of the ΛCDM run. This would make it easier to reach the threshold of reionisation at high reshifts in the ReBEL framework.

We further have to acknowledge that earlier formation of the DM haloes fostered by the ReBEL model, results in shifting a significant part of the violent merger events to higher redshifts. This indicates that there are many more haloes with extended steady accretion histories than expected in the standard ΛCDM structure formation scenarios. This would help thin-disk dominated galaxies to preserve their disks at low redshifts and would also imply that the exceptionally long period of peace experienced by the Milky-Way is no more so extraordinary (Hammer et al. 2007).

Finally, we have also shown that long-range scalar interactions between DM particles increase to some extent the masses of DM haloes. That is also an important effect that may help to understand the existence of objects like massive superclusters at moderate redshifts. The simulations presented in this *Letter* however do not encompass a large enough volume to study the formation of these massive objects in detail.

In conclusion, we have shown that the ReBEL model allows to alleviate some of the tension between predictions of the ΛCDM and observations (Benson et al. 2006; Kogut et al. 2003; Driver 2010; Stewart et al. 2008; Yamila Yaryura et al. 2010), however, as we only focused on the issue of formation time, it still needs to be investigated if and how internal properties of dark matter haloes are affected by the ReBEL formalism. One interesting quantity is the

halo concentration parameter, as it correlated with the halo formation time (eg. Wechsler et al. 2002). Preliminary work shows this to indeed be the case, with the mean concentration $\langle c_{vir} \rangle$ being 9 (11, 11, 13) for ΛCDM (*b05rs05*, *b05rs1*, *b1rs1*) for haloes with masses $M \sim 2 \times 10^{12} h^{-1} M_{\odot}$, respectively. However the issue of the halo concentration parameters require deeper studies and, since a more detailed study of the haloes internal properties is in progress, we will cover the concentration parameter issue in a forth-coming work.

## ACKNOWLEDGEMENTS

WAH would like to acknowledge the hospitality of the Departamento de Física Teórica at the Universidad Autónoma de Madrid during his short stay there. Some of the simulations used in this research was performed on the `halo` cluster at the Warsaw University Interdisciplinary Center for Mathematical and computational Modelling. WAH is supported by the Polish Ministry of Science grants no. NN203 394234 and NN203 386037. SRK acknowledges support by the Ministerio de Ciencia e Innovacion (MICINN) under the Consolider-Ingenio, SyeC project CSD-2007-00050. AK is supported by the MICINN through the Ramon y Cajal programme.

## REFERENCES

Baldi M., 2009, Nuclear Physics B Proceedings Supplements, 194, 178
Baldi M., Pettorino V., Robbers G., Springel V., 2010, MNRAS, 403, 1684
Benson A. J., Sugiyama N., Nusser A., Lacey C. G., 2006, MNRAS, 369, 1055
Bouwens R. J., Illingworth G. D., Labbe I., Oesch P. A., Carollo M., Trenti M., van Dokkum P. G., Franx M., Stiavelli M., Gonzalez V., Magee D., 2009, ArXiv e-prints
Bouwens R. J., Illingworth G. D., Oesch P. A., Stiavelli M., van Dokkum P., Trenti M., Magee D., Labbé I., Franx M., Carollo C. M., Gonzalez V., 2010, ApJ, 709, L133
Bower R. G., Benson A. J., Malbon R., Helly J. C., Frenk C. S., Baugh C. M., Cole S., Lacey C. G., 2006, MNRAS, 370, 645
Cen R., 2003, ApJ, 591, 12
Cen R., 2006, New Astronomy Review, 50, 191
Colless M., et al. 2003, ArXiv Astrophysics e-prints
Cowie L. L., Barger A. J., Bautz M. W., Brandt W. N., Garmire G. P., 2003, ApJ, 584, L57
Driver S. P., 2010, in V. P. Debattista & C. C. Popescu ed., American Institute of Physics Conference Series Vol. 1240 of American Institute of Physics Conference Series, The Decade of Galaxy Formation: Pitfalls in the Path Ahead. pp 17–28
Farrar G. R., Peebles P. J. E., 2004, Astrophys. J., 604, 1
Farrar G. R., Rosen R. A., 2007, Phys. Rev. Lett., 98, 171302
Fontanot F., De Lucia G., Monaco P., Somerville R. S., Santini P., 2009, MNRAS, 397, 1776
Gill S. P. D., Knebe A., Gibson B. K., 2004, MNRAS, 351, 399
González V., Labbé I., Bouwens R. J., Illingworth G., Franx M., Kriek M., Brammer G. B., 2010, ApJ, 713, 115
Governato F., Brook C. B., Brooks A. M., Mayer L., Willman B., Jonsson P., Stilp A. M., Pope L., Christensen C., Wadsley J., Quinn T., 2009, MNRAS, 398, 312
Gubser S. S., Peebles P. J. E., 2004a, Phys. Rev. D, 70, 123511
Gubser S. S., Peebles P. J. E., 2004b, Phys. Rev. D, 70, 123510
Haiman Z., 2004, ApJ, 613, 36
Hammer F., Puech M., Chemin L., Flores H., Lehnert M. D., 2007, ApJ, 662, 322
Hellwing W. A., 2010, Annalen der Physik, 19, 351
Hellwing W. A., Juszkiewicz R., 2009, Phys. Rev. D, 80, 083522
Hellwing W. A., Juszkiewicz R., van de Weygaert R., 2010, ArXiv e-prints
Keselman J. A., Nusser A., Peebles P. J. E., 2010, Phys. Rev. D, 81, 063521
Knollmann S. R., Knebe A., 2009, ApJS, 182, 608
Kogut A., Spergel D. N., Barnes C., Bennett C. L., Halpern M., Hinshaw G., Jarosik N., Limon M., Meyer S. S., Page L., Tucker G. S., Wollack E., Wright E. L., 2003, ApJS, 148, 161
Komatsu E., et al. 2010, ArXiv e-prints
Lacey C., Cole S., 1993, MNRAS, 262, 627
Lee J., 2010, ArXiv e-prints
Li B., Barrow J. D., 2010, ArXiv e-prints
Li B., Zhao H., 2010, Phys. Rev. D, 81, 104047
Lin W. P., Jing Y. P., Lin L., 2003, MNRAS, 344, 1327
Moster B. P., Macciò A. V., Somerville R. S., Johansson P. H., Naab T., 2010, MNRAS, 403, 1009
Nusser A., Gubser S. S., Peebles P. J., 2005, Phys. Rev. D, 71, 083505
Ouchi M., Mobasher B., Shimasaku K., Ferguson H. C., Fall S. M., Ono Y., Kashikawa N., Morokuma T., Nakajima K., Okamura S., Dickinson M., Giavalisco M., Ohta K., 2009, ApJ, 706, 1136
Peebles P. J. E., Nusser A., 2010, Nature, 465, 565
Springel V., 2005, MNRAS, 364, 1105
Stewart K. R., Bullock J. S., Wechsler R. H., Maller A. H., Zentner A. R., 2008, ApJ, 683, 597
Swinbank A. M., et al. 2007, MNRAS, 379, 1343
Tasitsiomi A., Kravtsov A. V., Gottlöber S., Klypin A. A., 2004, ApJ, 607, 125
Volonteri M., Haardt F., Madau P., 2003, ApJ, 582, 559
Wechsler R. H., Bullock J. S., Primack J. R., Kravtsov A. V., Dekel A., 2002, Astrophys. J., 568, 52
Willott C. J., Delorme P., Reylé C., Albert L., Bergeron J., Crampton D., Delfosse X., Forveille T., Hutchings J. B., McLure R. J., Omont A., Schade D., 2010, AJ, 139, 906
Yamila Yaryura C., Baugh C. M., Angulo R. E., 2010, ArXiv e-prints

This paper has been typeset from a TEX/ LATEX file prepared by the author.